



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.


MECHANICS.

196. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

From a uniform, solid right circular cone two planes cut off a portion such that the sections are similar ellipses with co-planar axes (not parallel). The centers of the elliptic faces are O_1 , O_2 , and the center of gravity of the solid is G . GX parallel to O_1O_2 cuts the axis of the cone in X . Find GX/O_1O_2 in terms of the ratio of the major axes of the ellipses.

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let ABC be the cone; BR , DF the major axes of the required ellipses. For calculations we will only consider the cone ADE and the plane DF .



Let $x^2 + z^2 = n^2(c-y)^2$ be the cone, $x = my - R$ the plane, where $R = DL$, $c = AL$, $h = FJ$, $r = \frac{1}{2}GF$, $n = \tan LAE = (R-r)/h$, $m = \tan FDE = (R+r)/h$, $c = Rh/(R-r)$. Let $my - R = x_1$, $n(c-y) = x_2$, $(x+nc)/n = y_1$, $(x+R)/m = y_2$, $(nc-x)/n = y_3$.

$$\begin{aligned} \therefore v &= 2 \int_0^h \int_{-x_2}^{x_1} \sqrt{[n^2(c-y)^2 - x^2]} dy dx + 2 \int_h^c \int_{-x_2}^{x_2} \sqrt{[n^2(c-y)^2 - x^2]} dy dx \\ &= \int_0^h \left\{ n^2(c-y)^2 \left[\frac{1}{2}\pi + \sin^{-1} \left(\frac{my-R}{n(c-y)} \right) \right] \right. \\ &\quad \left. + (my-R) \sqrt{[n^2(c-y)^2 - (my-R)^2]} \right\} dy + \pi n^2 \int_h^c (c-y)^2 dy = \frac{\frac{1}{3}\pi R r h \sqrt{Rr}}{R-r}. \end{aligned}$$

$$\therefore v = \frac{1}{3}\pi (AD \cdot AF)^{\frac{2}{3}} \sin^2 \beta \cos \beta, \text{ where } \angle LAE = \beta.$$

$$\begin{aligned} \bar{x}v &= 2 \int_0^h \int_{-x_2}^{x_1} x \sqrt{[n^2(c-y)^2 - x^2]} dy dx + 2 \int_h^c \int_{-x_2}^{x_2} x \sqrt{[n^2(c-y)^2 - x^2]} dy dx \\ &= -\frac{2}{3} \int_0^h [n^2(c-y)^2 - (my-R)^2]^{\frac{3}{2}} dy \\ &= -\frac{2}{3(m^2 - n^2)^{\frac{5}{2}}} \int_{-u_1}^{u_1} [(ncm - Rn)^2 - u^2]^{\frac{3}{2}} du \end{aligned}$$

where $u = (m^2 - n^2)y + n^2c - Rm$, $u_1 = 2rR/h$.

$$\therefore \bar{x}v = -\frac{1}{3}\pi R r h \sqrt{Rr} = -\frac{1}{3}\pi (AD - AF) (AD \cdot AF)^{\frac{2}{3}} \sin^3 \beta \cos \beta.$$

$$\begin{aligned} \bar{y}v &= 2 \int_0^h \int_{-x_2}^{x_1} y \sqrt{[n^2(c-y)^2 - x^2]} dy dx + 2 \int_h^c \int_{-x_2}^{x_2} y \sqrt{[n^2(c-y)^2 - x^2]} dy dx \\ &= 2c \int_0^h \int_{-x_2}^{x_1} \sqrt{[n^2(c-y)^2 - x^2]} dy dx + 2c \int_h^c \int_{-x_2}^{x_2} \sqrt{[n^2(c-y)^2 - x^2]} dy dx \\ &\quad - 2 \int_{-r}^r \int_{y_2}^{y_3} (c-y) \sqrt{[n^2(c-y)^2 - x^2]} dx dy \\ &\quad - 2 \int_{-R}^{-r} \int_{y_2}^{y_1} (c-y) \sqrt{[n^2(c-y)^2 - x^2]} dx dy \\ &\quad - 2 \int_{-r}^r \int_h^{y_3} (c-y) \sqrt{[n^2(c-y)^2 - x^2]} dx dy \\ &= \frac{\frac{1}{3}\pi c R r h \sqrt{Rr}}{R-r} - \frac{2}{3n^2 m^3} \int_{-R}^r [n^2(cm - R - x)^2 - m^2 x^2]^{\frac{3}{2}} dx = \frac{\frac{1}{3}\pi c R r h \sqrt{Rr}}{R-r} \end{aligned}$$

$$\frac{2}{3n^2m^3(m^2-n^2)^{\frac{3}{2}}} \int_{-u_2}^{u_1} [(ncm^2-nRm)^2-x^2]^{\frac{1}{2}} du,$$

where $u=(m^2-n^2)x^2+cmn^2-Rn^2$, $u_2=\frac{2Rr(R+r)}{h^2}$.

$$\therefore \bar{y}v = \frac{\pi R r h^2 \sqrt{(Rr)(5R-3r)}}{24(R-r)^2} = \frac{\pi}{24} (5AD-3AF)(AD.AF)^{\frac{3}{2}} \sin^2\beta \cos^2\beta.$$

\bar{x} , \bar{y} give the center of gravity of the cone ADF . The center of gravity G of the portion $FDBRF$ is given as follows:

$$\bar{x}_1 = \frac{3[(AD-AF)(AD.AF)^{\frac{3}{2}} - (AB-AR)(AB.AR)^{\frac{3}{2}}] \sin\beta}{8[(AB.AR)^{\frac{3}{2}} - (AD.AF)^{\frac{3}{2}}]} = a.$$

$$\bar{y}_1 = \frac{[(5AB-3AR)(AB.AR)^{\frac{3}{2}} - (5AD-3AF)(AD.AF)^{\frac{3}{2}}] \cos\beta}{8[(AB.AR)^{\frac{3}{2}} - (AD.AF)^{\frac{3}{2}}]} = b.$$

Now $DF = \sqrt{(AD^2 + AF^2 - 2AD.AF \cos 2\beta)}$,

$BR = \sqrt{(AB^2 + AR^2 - 2AB.AR \cos 2\beta)}$.

Coordinates of O_1 are, $[AB - \frac{1}{2}(AD+AF)] \cos\beta$, and, $\frac{1}{2}(AD-AF) \sin\beta$,

Coordinates of O_2 are, $\frac{1}{2}(AB-AR) \cos\beta$, and, $\frac{1}{2}(AB-AR) \sin\beta$.

$\therefore O_1O_2 = \frac{1}{2}[(AB-AD)^2 + (AR-AF)^2 + 2(AB-AD)(AR-AF) \cos 2\beta]^{\frac{1}{2}}$.

GX makes with the axis of abscissas an angle whose tangent is

$$\left(\frac{AD+AF-AB-AR}{AB+AF-AR-AD} \right) \cot\beta = \mu.$$

$\therefore y-b = \mu(x-a)$ is the equation to GX . This meets the axis of ordinates in $x=0$, $y=b-\mu a$.

$\therefore GX = a\sqrt{1+\mu^2}$

$$= \frac{a\sqrt{[(AB-AD)^2 + (AR-AF)^2 + 2(AB-AD)(AR-AF) \cos 2\beta]}}{(AB+AF-AR-AD) \sin\beta}$$

$$= \frac{2aO_1O_2}{(AB+AF-AR-AD) \sin\beta}$$

$$\therefore \frac{GX}{O_1O_2} = \frac{2a}{(AB+AF-AR-AD) \sin\beta}$$

$$= \frac{6[(AD-AF)(AD.AF)^{\frac{3}{2}} - (AB-AR)(AB.AR)^{\frac{3}{2}}]}{8(AB+AF-AR-AD)[(AB.AR)^{\frac{3}{2}} - (AD.AF)^{\frac{3}{2}}]}.$$

Let $\angle FDE = \theta$, $\angle RBC = \phi$, semi-minor axis of ellipse $DF = p$, of ellipse $BR = q$. Then

$$p = \sqrt{(HO_1 \cdot O_1 K)} = \sqrt{(AD \cdot AF)} \sin \beta, \quad q = \sqrt{(ZO_2 \cdot O_2 I)} = \sqrt{(AB \cdot AR)} \sin \beta,$$

$$AD - AF = FE = \frac{DF \sin \theta}{\cos \beta}, \quad AB - AR = RC = \frac{BR \sin \phi}{\cos \beta}.$$

$$\therefore \frac{GX}{O_1 O_2} = \frac{6(p^3 DF \sin \theta - q^3 BR \sin \phi)}{8(BR \sin \phi - DF \sin \theta)(q^3 - p^3)}.$$

Let $DF = 2P$, $BR = 2Q$. Then

$$p = \frac{P \sqrt{[\cos(\theta + \beta) \cos(\theta - \beta)]}}{\cos \beta} = \frac{P \sqrt{[\cos^2 \theta - \sin^2 \beta]}}{\cos \beta}, \quad q = \frac{Q \sqrt{[\cos^2 \theta - \sin^2 \beta]}}{\cos \beta}.$$

$$\therefore \frac{GX}{O_1 O_2} = \frac{6[P^4 \sin \theta (\cos^2 \theta - \sin^2 \beta)^{\frac{3}{2}} - Q^4 \sin \phi (\cos^2 \phi - \sin^2 \beta)^{\frac{3}{2}}]}{8\{[Q \sin \phi - P \sin \theta][Q^3 (\cos^2 \phi - \sin^2 \beta)^{\frac{3}{2}} - P^3 (\cos^2 \theta - \sin^2 \beta)^{\frac{3}{2}}]\}}.$$

If the planes of the ellipses are parallel, $\theta = \phi$, and we get

$$\frac{GX}{O_1 O_2} = \frac{6(P^4 - Q^4)}{8(Q - P)(Q^3 - P^3)} = \frac{6(P^2 + Q^2)(P + Q)}{P^3 - Q^3}.$$

MISCELLANEOUS.

165. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

$$\text{Prove that } \tan^{-1} \frac{n}{n+1} + \tan^{-1} \frac{1}{2n+1} = \frac{1}{2} \pi.$$

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.; A. H. HOLMES, Brunswick, Me.; J. EDWARD SANDERS, Reinersville, O.; FRANK M. DRYZER, A. B., Knoxville, Tenn.; and PROF. J. W. NICHOLSON, State University, La.

$$\text{Let } \alpha = \tan^{-1} \frac{n}{n+1} \text{ and } \beta = \tan^{-1} \frac{1}{2n+1}.$$

$$\text{Then } \tan \alpha = \frac{n}{n+1}, \tan \beta = \frac{1}{2n+1}, \text{ and } \tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$$

$$= \frac{\frac{n}{n+1} + \frac{1}{2n+1}}{1 - \frac{n}{(n+1)(2n+1)}} = \frac{2n^2 + 2n + 1}{2n^2 + 2n + 1} = 1.$$

$$\therefore \alpha + \beta = \frac{1}{2} \pi.$$

Also solved by G. W. Greenwood.

166. Proposed by F. H. SAFFORD, Ph. D., The University of Pennsylvania.

Several equal rectangular boxes are placed in a row with uniform intervals between the boxes and a passageway along one side of the row. Find the least width of the passageway permitting a box to be removed from the row without moving adjacent boxes. This problem arose during the construction of a room for storage batteries.